

Erratum to: “Nuclear Effects on $R = \sigma_L/\sigma_T$ in Deep-Inelastic Scattering”
Phys. Lett. B 475 (2000) 386

A. Airapetian,³¹ N. Akopov,³¹ Z. Akopov,³¹ M. Amarian,^{26,31} V.V. Ammosov,²⁴ E.C. Aschenauer,⁶ R. Avakian,³¹
 A. Avetissian,³¹ E. Avetissian,³¹ P. Bailey,¹⁵ V. Baturin,²³ C. Baumgarten,²¹ M. Beckmann,⁵ S. Belostotski,²³
 S. Bernreuther,²⁹ N. Bianchi,¹⁰ H.P. Blok,^{22,30} H. Böttcher,⁶ A. Borissov,¹⁹ O. Bouhali,²² M. Bouwhuis,¹⁵
 J. Brack,⁴ S. Brauksiepe,¹¹ A. Brüll,¹⁸ I. Brunn,⁸ G.P. Capitani,¹⁰ H.C. Chiang,¹⁵ G. Ciullo,⁹ G.R. Court,¹⁶
 P.F. Dalpiaz,⁹ R. De Leo,³ L. De Nardo,¹ E. De Sanctis,¹⁰ E. Devitsin,²⁰ P.K.A. de Witt Huberts,²² P. Di Nezza,¹⁰
 M. Düren,¹³ M. Ehrenfried,⁶ A. Elalaoui-Moulay,² G. Elbakian,³¹ F. Ellinghaus,⁶ U. Elschenbroich,¹¹ J. Ely,⁴
 R. Fabbri,⁹ A. Fantoni,¹⁰ A. Fechtchenko,⁷ L. Felawka,²⁸ H. Fischer,¹¹ B. Fox,⁴ J. Franz,¹¹ S. Frullani,²⁶
 Y. Gärber,⁸ G. Gapienko,²⁴ V. Gapienko,²⁴ F. Garibaldi,²⁶ E. Garutti,²² G. Gavrilov,²³ V. Gharibyan,³¹
 G. Graw,²¹ O. Grebenioun,²³ P.W. Green,^{1,28} L.G. Greeniaus,^{1,28} A. Gute,⁸ W. Haeberli,¹⁷ K. Hafidi,²
 M. Hartig,²⁸ D. Hasch,¹⁰ D. Heesbeen,²² F.H. Heinsius,¹¹ M. Henoch,⁸ R. Hertenberger,²¹ W.H.A. Hesselink,^{22,30}
 Y. Holler,⁵ B. Hommez,¹² G. Iarygin,⁷ A. Izotov,²³ H.E. Jackson,² A. Jgoun,²³ R. Kaiser,¹⁴ E. Kinney,⁴
 A. Kisselev,²³ P. Kitching,¹ K. Königsmann,¹¹ H. Kolster,¹⁸ M. Kopytin,²³ V. Korotkov,⁶ E. Kotik,¹ V. Kozlov,²⁰
 B. Krauss,⁸ V.G. Krivokhijine,⁷ L. Lagamba,³ L. Lapikás,²² A. Laziev,^{22,30} P. Lenisa,⁹ P. Liebing,⁶
 T. Lindemann,⁵ W. Lorenzon,¹⁹ N.C.R. Makins,¹⁵ H. Marukyan,³¹ F. Masoli,⁹ F. Menden,¹¹ V. Mexner,²²
 N. Meyners,⁵ O. Mikloukho,²³ C.A. Miller,^{1,28} V. Muccifora,¹⁰ A. Nagaitsev,⁷ E. Nappi,³ Y. Naryshkin,²³
 A. Nass,⁸ K. Negodaeva,⁶ W.-D. Nowak,⁶ K. Oganessyan,^{5,10} H. Ohsuga,²⁹ G. Orlandi,²⁶ S. Podiathev,⁸
 S. Potashov,²⁰ D.H. Potterveld,² M. Raithel,⁸ D. Reggiani,⁹ P.E. Reimer,² A. Reischl,²² A.R. Reolon,¹⁰ K. Rith,⁸
 G. Rosner,¹⁴ A. Rostomyan,³¹ D. Ryckbosch,¹² Y. Sakemi,²⁹ I. Sanjiev,^{2,23} F. Sato,²⁹ I. Savin,⁷ C. Scarlett,¹⁹
 A. Schäfer,²⁵ C. Schill,¹¹ G. Schnell,⁶ K.P. Schüller,⁵ A. Schwind,⁶ J. Seibert,¹¹ B. Seitz,¹ R. Shanidze,⁸
 T.-A. Shibata,²⁹ V. Shutov,⁷ M.C. Simani,^{22,30} K. Sinram,⁵ M. Stancari,⁹ M. Statera,⁹ E. Steffens,⁸
 J.J.M. Steijger,²² J. Stewart,⁶ U. Stösslein,⁴ K. Suetsugu,²⁹ H. Tanaka,²⁹ S. Taroian,³¹ A. Terkulov,²⁰
 S. Tessarin,⁹ E. Thomas,¹⁰ A. Tkabladze,⁶ M. Tytgat,¹² G.M. Urciuoli,²⁶ G. van der Steenhoven,²²
 R. van de Vyver,¹² M.C. Vetterli,^{27,28} V. Vikhrov,²³ M.G. Vincet,¹ J. Visser,²² J. Volmer,⁶ C. Weiskopf,⁸
 J. Wendland,^{27,28} J. Wilbert,⁸ T. Wise,¹⁷ S. Yen,²⁸ S. Yoneyama,²⁹ B. Zihlmann,^{22,30} and H. Zohrabian³¹

(The HERMES Collaboration)

¹Department of Physics, University of Alberta, Edmonton, Alberta T6G 2J1, Canada

²Physics Division, Argonne National Laboratory, Argonne, Illinois 60439-4843, USA

³Istituto Nazionale di Fisica Nucleare, Sezione di Bari, 70124 Bari, Italy

⁴Nuclear Physics Laboratory, University of Colorado, Boulder, Colorado 80309-0446, USA

⁵DESY, Deutsches Elektronen-Synchrotron, 22603 Hamburg, Germany

⁶DESY Zeuthen, 15738 Zeuthen, Germany

⁷Joint Institute for Nuclear Research, 141980 Dubna, Russia

⁸Physikalisches Institut, Universität Erlangen-Nürnberg, 91058 Erlangen, Germany

⁹Istituto Nazionale di Fisica Nucleare, Sezione di Ferrara and

Dipartimento di Fisica, Università di Ferrara, 44100 Ferrara, Italy

¹⁰Istituto Nazionale di Fisica Nucleare, Laboratori Nazionali di Frascati, 00044 Frascati, Italy

¹¹Fakultät für Physik, Universität Freiburg, 79104 Freiburg, Germany

¹²Department of Subatomic and Radiation Physics, University of Gent, 9000 Gent, Belgium

¹³Physikalisches Institut, Universität Gießen, 35392 Gießen, Germany

¹⁴Department of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, United Kingdom

¹⁵Department of Physics, University of Illinois, Urbana, Illinois 61801, USA

¹⁶Physics Department, University of Liverpool, Liverpool L69 7ZE, United Kingdom

¹⁷Department of Physics, University of Wisconsin-Madison, Madison, Wisconsin 53706, USA

¹⁸Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

¹⁹Randall Laboratory of Physics, University of Michigan, Ann Arbor, Michigan 48109-1120, USA

²⁰Lebedev Physical Institute, 117924 Moscow, Russia

²¹Sektion Physik, Universität München, 85748 Garching, Germany

²²Nationaal Instituut voor Kernfysica en Hoge-Energiefysica (NIKHEF), 1009 DB Amsterdam, The Netherlands

²³Petersburg Nuclear Physics Institute, St. Petersburg, Gatchina, 188350 Russia

²⁴Institute for High Energy Physics, Protvino, Moscow oblast, 142284 Russia

²⁵Institut für Theoretische Physik, Universität Regensburg, 93040 Regensburg, Germany

²⁶Istituto Nazionale di Fisica Nucleare, Sezione Roma 1, Gruppo Sanità

and Physics Laboratory, Istituto Superiore di Sanità, 00161 Roma, Italy

²⁷Department of Physics, Simon Fraser University, Burnaby, British Columbia V5A 1S6, Canada

²⁸TRIUMF, Vancouver, British Columbia V6T 2A3, Canada

²⁹Department of Physics, Tokyo Institute of Technology, Tokyo 152, Japan

³⁰Department of Physics and Astronomy, Vrije Universiteit, 1081 HV Amsterdam, The Netherlands

³¹Yerevan Physics Institute, 375036 Yerevan, Armenia

(Dated: February 7, 2008)

This erratum revokes the main conclusion of a Letter that reported measurements of cross sections for deep-inelastic scattering (DIS) of leptons on ^3He and ^{14}N targets, expressed as ratios of σ_A/σ_D to the cross section on the deuterium target [*]. In the particular kinematic domain $x < 0.03$ with $Q^2 < 1.25 \text{ GeV}^2$, σ_A/σ_D was reported to differ as much as 35% from earlier such measurements at higher energies. As the only significant difference from the earlier measurements appeared to be the kinematic variable y , and hence the polarization parameter ϵ , the new results were interpreted as evidence for a nuclear influence on the ratio R of the cross sections for longitudinal and transverse photons. This anomaly has now been discovered to be due to a peculiar instrumental effect.

Radiative corrections do not cancel in the ratio σ_A/σ_D because the yield of radiative events associated with elastic scattering scales nonlinearly, with Z^2 . At small values of *apparent* x and Q^2 (inferred from the measured angle and energy of the scattered lepton), corresponding to large values of y , the contribution from radiative elastic scattering becomes large. Unlike radiation associated with inelastic processes, which is predominantly emitted in the direction of either the beam lepton (initial state radiation or ISR) or the scattered lepton (final state radiation or FSR), the hard photons associated with nuclear elastic scattering involve negligible momentum transfer q to the target nucleus (Compton peak). There are two reasons for this. One is that the Bethe-Heitler cross section for radiative elastic processes predicts that in kinematic conditions corresponding to quite small values of *apparent* x and Q^2 , the Compton peak becomes much more prominent compared to ISR and FSR, simply because smaller values of q become kinematically available, and the cross section is modulated by a factor of $1/q^4$. This is illustrated in Fig. 1, which shows the nuclear-elastic Bethe-Heitler cross section in two different coplanar kinematic situations, both with and without including the nuclear form factor. This latter comparison reveals the other reason — that the nuclear form factor strongly suppresses the cross section for significant momentum transfer to the target, leaving only the Compton peak.

With negligible nuclear recoil momentum, essentially all of the transverse momentum of the scattered lepton must be balanced by that of the radiated hard photon, which also carries away most of the beam energy at these large values of *apparent* y . Hence we have

$$(1 - y) \sin \theta_{e'} = y \sin \theta_\gamma, \quad (1)$$




FIG. 1: The nuclear-elastic Bethe-Heitler cross section [2] on ^{14}N for two different coplanar kinematic conditions as labelled in terms of *apparent* DIS kinematic variables. The continuous curves include the effects of the nuclear form factor.

showing that at large y , the angle of the high-energy photon on the opposite side of the beam line is correspondingly smaller than that of the scattered lepton, but not negligible. In the mirror-symmetric open geometry of the HERMES spectrometer, this can have drastic consequences. These energetic photons from nuclear targets have a high probability of hitting the detector frames surrounding the beam line in front of the dipole magnet, and producing extensive electromagnetic showers that cause very high hit multiplicities in these tracking detectors. For many of these nuclear-elastic events, track reconstruction is therefore impossible, resulting in a large tracking inefficiency that is strictly correlated with only this process and kinematic situation.

This problem is pernicious because it is far from apparent in the experimental data. The event trigger rate for real DIS events is typically very small compared to that from hadron background. Only after event reconstruction can all of the particle identification criteria be

applied to eliminate the hadrons. However, event reconstruction is impossible for the affected radiative elastic events, so they remain hidden in the dominant hadron background and lost to the analysis, even though they are included in the radiative corrections. A simulation of the experiment reveals the problem only if it includes both the nuclear target with its particular radiative effects, and a complete treatment of showers in material *outside* of the geometric acceptance. This was not included in the data analysis for the original Letter. Subsequent such studies [1] revealed that these showers can indeed account for the differences between the HERMES results and previous measurements in similar kinematic regions. The strong correlation of the shower phenomenon with the kinematic variable y accounts for its correlation also with the photon polarization parameter ϵ , which led to the incorrect interpretation in terms of R_A/R_D . This interpretation became untenable when analyzing $^{84}\text{Kr}/^2\text{H}$ cross section ratios extracted from data collected after the publication of the $^{14}\text{N}/^2\text{H}$ data: for this heavy target nucleus the instrumental effects described above are so severe that the extracted Born cross section ratio became unphysical. The corrected $^3\text{He}/^2\text{H}$ and $^{14}\text{N}/^2\text{H}$ data and the new $^{84}\text{Kr}/^2\text{H}$ data are presented in the accompanying

paper [1].

The part of the HERMES kinematic region affected by the correlated background from nuclear targets is restricted to $x < 0.06$ with $Q^2 < 2 \text{ GeV}^2$. Polarized DIS data from hydrogen, deuterium and helium-3 targets are unaffected by this effect, because of both the more restricted kinematic range, and the much smaller value of Z^2 modulating the elastic Bethe-Heitler cross section. Semi-inclusive data are also unaffected even with nuclear targets [3], as radiative elastic events are excluded by the presence of a hadron in the final state.

-
- [*] All symbols used here are defined in the original Letter.
 [1] A. Airapetian *et al.* (HERMES), Phys. Lett. B, this volume.
 [2] C. Itzykson and J.-B. Zuber, *Quantum Field Theory*, New York, McGraw-Hill, 1985 (Eq. 5.150)
 [3] K. Ackerstaff *et al.* (HERMES), Phys. Rev. Lett. **82** (1999) 3025;
 A. Airapetian *et al.* (HERMES), Eur. Phys. J. C **20** (2001) 479.